

Original article

Influence of birth cohort on age of onset cluster analysis in bipolar I disorder

M. Bauer [a,*](#), T. Glenn [b](#), M. Alda [c](#), O.A. Andreassen [d](#), E. Angelopoulos [e](#), R. Arda [f](#), C. Baethge [g](#), R. Bauer [a](#), F. Bellivier [h,aw](#), R.H. Belmaker [i](#), M. Berk [j,k](#), T.D. Bjella [d](#), L. Bossini [l](#), Y. Bersudsky [i](#), E.Y.W. Cheung [m](#), J. Conell [a](#), M. Del Zompo [n](#), S. Dodd [j,o](#), B. Etain [p](#), A. Fagiolini [l](#), M.A. Frye [q](#), K.N. Fountoulakis [r](#), J. Garneau-Fournier [s](#), A. Gonzalez-Pinto [t](#), H. Harima [u](#), S. Hassel [v](#), C. Henry [p](#), A. Iacovides [r](#), E.T. Isometsa [w,x](#), F. Kapczinski [y](#), S. Kliwicki [z](#), B. Koenig [aa](#), R. Krogh [ab](#), M. Kunz [y](#), B. Lafer [ac](#), E.R. Larsen [ab](#), U. Lewitzka [a](#), C. Lopez-Jaramillo [ad](#), G. MacQueen [v](#), M. Manchia [c](#), W. Marsh [ae](#), M. Martinez-Cengotitabengoa [t](#), I. Melle [d](#), S. Monteith [af](#), G. Morken [ag](#), R. Munoz [ah](#), F.G. Nery [ac](#), C. O'Donovan [c](#), Y. Osher [i](#), A. Pfennig [a](#), D. Quiroz [ai](#), R. Ramesar [aj](#), N. Rasgon [s](#), A. Reif [ak](#), P. Ritter [a](#), J.K. Rybakowski [z](#), K. Sagduyu [al](#), A.M. Scippa [am](#), E. Severus [a](#), C. Simhandl [aa](#), D.J. Stein [an](#), S. Strejilevich [ao](#), A. Hatim Sulaiman [ap](#), K. Suominen [aq](#), H. Tagata [u](#), Y. Tatebayashi [ar](#), C. Torrent [as](#), E. Vieta [as](#), B. Viswanath [at](#), M.J. Wanchoo [q](#), M. Zetin [au](#), P.C. Whybrow [av](#)

[a](#) Department of Psychiatry and Psychotherapy, Universitätsklinikum Carl Gustav Carus, Technische Universität Dresden, Germany Fetscherstr. 74, 01307 Dresden, Germany [b](#) ChronoRecord Association, Fullerton, CA, USA [c](#) Department of Psychiatry, Dalhousie University, Halifax, NS, Canada

[d](#) NORMENT - K.G. Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, Oslo, Norway [e](#)

Department of Psychiatry, University of Athens Medical School, Eginition Hospital, Athens, Greece [f](#) Unit of Clinical Pharmacology, University Hospital of Cagliari, Cagliari, Italy

[g](#) Department of Psychiatry and Psychotherapy, University of Cologne Medical School, Cologne, Germany [h](#) Psychiatrie, GH Saint-Louis – Lariboisière – F.-Widal, AP-HP, INSERM UMR-S1144, Faculté de Médecine, Université D.-Diderot, Paris, France [i](#)

Department of Psychiatry, Faculty of Health Sciences, Ben Gurion University of the Negev, Beer Sheva Mental Health Center, Beer Sheva, Israel [j](#) IMPACT Strategic Research Centre, School of Medicine, Deakin University, Geelong, Victoria 3220, Australia [k](#) Department of Psychiatry, ORYGEN Youth Health Research Centre, Centre for Youth Mental Health and the Florey Institute for Neuroscience and Mental Health, University of Melbourne, Parkville, Victoria 3052, Australia [l](#) Department of Molecular Medicine and Department of Mental Health (DAI), University of Siena and University of Siena Medical Center (AOUS), Siena, Italy [m](#) Department of General Adult Psychiatry, Castle Peak Hospital, Hong Kong [n](#) Section of Neurosciences and Clinical Pharmacology, Department of Biomedical Sciences, University of Cagliari, Sardinia, Italy

[o](#) Department of Psychiatry, University of Melbourne, Parkville, Victoria 3052, Australia [p](#) AP-HP, Hôpitaux Universitaires Henri-Mondor, INSERM U955 (IMRB), Université Paris Est, and FondaMental Fondation, Créteil, France [q](#) Department of Psychiatry &

Psychology, Mayo Clinic Depression Center, Mayo Clinic, Rochester, MN, USA [r](#) 3rd Department of Psychiatry, Division of Neurosciences, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece [s](#) Department of Psychiatry and Behavioral Sciences, Stanford School of Medicine, Palo Alto, CA, USA [t](#) Department of Psychiatry, University Hospital of Alava, University of the Basque Country, CIBERSAM, Vitoria, Spain [u](#) Department of Psychiatry, Tokyo Metropolitan Matsuzawa Hospital, Setagaya, Tokyo, Japan [v](#) Department of Psychiatry, Faculty of Medicine, University of Calgary, Calgary, AB, Canada [w](#) Department of Psychiatry, Institute of Clinical Medicine, University of Helsinki, Helsinki, Finland [x](#) National Institute for Health and Welfare,

Helsinki, Finland y Laboratory of Molecular Psychiatry, Hospital de Clínicas de Porto Alegre, Federal University of Rio Grande do Sul, Porto Alegre, Brazil z Department of Adult Psychiatry, Poznan University of Medical Sciences, Poznan, Poland aa BIPOLAR Zentrum Wiener Neustadt, Wiener Neustadt, Austria ab Department of Affective Disorders, Q, Mood Disorders Research Unit, Aarhus University Hospital, Aarhus, Denmark ac Bipolar Disorder Research Program, Department of Psychiatry, University of São Paulo Medical School, São Paulo, Brazil ad Mood Disorders Program, Fundación San Vicente de Paul, Department of Psychiatry, Universidad de Antioquia, Medellín, Colombia
 e Department of Psychiatry, University of Massachusetts, Worcester, MA, USA af Michigan State University College of Human Medicine, Traverse City Campus, Traverse City, MI, USA ag Department of Neuroscience, NTNU, and St Olavs' University Hospital, Trondheim, Norway ah Department of Psychiatry, University of California San Diego, San Diego, CA, USA ai Department of Psychiatry, Diego Portales University, Santiago, Chile aj UCT/MRC Human Genetics Research Unit, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa ak Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, Goethe-University Frankfurt am Main, Germany al Department of Psychiatry, University of Missouri Kansas City School of Medicine, Kansas City, MO, USA am Department of Neuroscience and Mental Health, Federal University of Bahia, Salvador, Brazil an Department of Psychiatry, University of Cape Town, Cape Town, South Africa ao Bipolar Disorder Program, Neuroscience Institute, Favaloro University, Buenos Aires, Argentina ap Department of Psychological Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia aq City of Helsinki, Department of Social Services and Health Care, Psychiatry, Helsinki, Finland ar Schizophrenia & Affective Disorders Research Project, Tokyo Metropolitan Institute of Medical Science, Seitama, Tokyo, Japan as Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain at Department of Psychiatry, NIMHANS, Bangalore 560029, India au Department of Psychology, Chapman University, Orange, CA, USA av Department of Psychiatry and Biobehavioral Sciences, Semel Institute for Neuroscience and Human Behavior University of California Los Angeles (UCLA), Los Angeles, CA, USA aw FondaMental Foundation, Creteil, France

Corresponding author. Tel.: +49 351 458 0; fax: +49 30 450 51 79 62. E-mail address: michael.bauer@uniklinikum-dresden.de (M. Bauer).

<http://dx.doi.org/10.1016/j.eurpsy.2014.10.005>

0924-9338/ß 2014 Elsevier Masson SAS. All rights reserved.

ABSTRACT

Purpose: Two common approaches to identify subgroups of patients with bipolar disorder are clustering methodology (mixture analysis) based on the age of onset, and a birth cohort analysis. This study investigates if a birth cohort effect will influence the results of clustering on the age of onset, using a large, international database.

Methods: The database includes 4037 patients with a diagnosis of bipolar I disorder, previously collected at 36 collection sites in 23 countries. Generalized estimating equations (GEE) were used to adjust the data for country median age, and in some models, birth cohort. Model-based clustering (mixture analysis) was then performed on the age of onset data using the residuals. Clinical variables in subgroups were compared.

Results: There was a strong birth cohort effect. Without adjusting for the birth cohort,

three subgroups were found by clustering. After adjusting for the birth cohort or when considering only those born after 1959, two subgroups were found. With results of either two or three subgroups, the youngest subgroup was more likely to have a family history of mood disorders and a first episode with depressed polarity. However, without adjusting for birth cohort (three subgroups), family history and polarity of the first episode could not be distinguished between the middle and oldest subgroups.

Conclusion: These results using international data confirm prior findings using single country data, that there are subgroups of bipolar I disorder based on the age of onset, and that there is a birth cohort effect. Including the birth cohort adjustment altered the number and characteristics of subgroups detected when clustering by age of onset. Further investigation is needed to determine if combining both approaches will identify subgroups that are more useful for research.

1. Introduction

The age of disease onset is often analyzed to identify patient subgroups that differ in clinical course or genetic profile. Two general approaches to grouping data from patients with bipolar disorder have provided important and replicated findings. The first approach uses a clustering methodology (mixture analysis) to determine the optimal number of distinct subgroups in a sample based on the age of onset distribution [8]. Using this clustering methodology, researchers have identified three onset subgroups, with the youngest subgroup having the most severe course of illness and highest likelihood of a family history of mood disorders [2,8,9,22,24,35,37]. The second approach groups the data in a sample by patient year of birth and analyzes for a birth cohort effect [21].

Researchers have detected a strong birth cohort effect in bipolar disorder, with successive generations experiencing an earlier age of onset [12,14,15,21,23,33,34]. The purpose of this analysis is to evaluate whether a birth cohort effect influences the results of clustering based on the age of onset using a large international database of patients with bipolar I disorder [4]. This is important because the birth cohort may modify the number and composition of subgroups, which in turn may affect the subsequent search for distinct and meaningful clinical and genetic profiles.

2. Methods

2.1. Data collection

The data in this analysis were collected for a study of the impact of solar insolation on the age of onset of bipolar disorder, and are described in detail elsewhere [4,5]. The diagnosis of bipolar disorder was made by a psychiatrist according to DSM-IV criteria. The patient data were obtained retrospectively at 36 collection sites in 23 countries. In 20 sites, data were obtained by a combination of direct interviews and record review, in 8 sites primarily by direct interview and in 8 sites by record review. The age of onset was defined as the first occurrence of an episode of depression, mania or hypomania according to DSM-IV criteria. Additional data included a family history of any mood disorder in a first degree relative, and the polarity of the first episode (depressed, manic or hypomanic). Study approval from institutional review boards was obtained according to local requirements.

Data from 5465 patients with bipolar disorder were obtained from 36 collection sites: Aarhus, Denmark (n = 66); Athens, Greece (n = 51); Bangalore, India (n = 99); Barcelona, Catalonia, Spain (n = 200); Beer Sheva, Israel (n = 105); Buenos Aires, Argentina (n = 95); Cagliari, Sardinia, Italy (n = 206); Calgary, Canada (n = 126); Cape Town, South Africa (n = 100); Dresden, Germany (n = 35); Halifax, Canada (n = 102); Helsinki, Finland (n = 191); Hong Kong (n = 50); Kansas City, KS, USA (n = 21); Kuala

Lumpur, Malaysia (n = 121); Los Angeles, CA, USA (n = 206); Medellin, Colombia (n = 189); Melbourne/Geelong, Australia (n = 161); Oslo, Norway (n = 127); Palo Alto, CA, USA (n = 48); Paris, France (n = 468); Porto Alegre, Brazil (n = 205); Poznan, Poland (n = 102); Rochester, MN, USA (n = 141); San Diego, CA, USA (n = 55); São Paulo, Brazil (n = 248); Salvador, Brazil (n = 121); Santiago, Chile (n = 346); Siena, Italy (n = 60); Thessaloniki, Greece (n = 52); Tokyo, Japan (n = 120); Trondheim, Norway (n = 238); Vitoria-Basque Country, Spain (n = 343); Worcester, MA, USA (n = 58); Wiener Neustadt, Germany (n = 356).

2.2. Database characteristics

Of the 5465 total patients 4037 were diagnosed with bipolar I disorder, 1236 with bipolar II and 192 with bipolar NOS. Due to a large imbalance in the diagnosis of bipolar I disorder at the collection sites, varying from 23% to 99%, only the 4037 patients with a diagnosis of bipolar I disorder were included in this analysis. Of the 4037 patients, 2374 (58.8%) were female and 1663 (41.2%) were male. Onset occurred in the southern hemisphere for 1043 (25.8%) of the patients.

The mean age of the 4037 patients was 48.1 \pm 14.5 years. The unadjusted mean age of onset for the 4037 patients was 25.4 years, similar to 25.7 years (n = 1665) in other research [3]. Family history was available for 3334 (82.6%) of the 4037 patients. Of the 3334 patients, 1848 (55.4%) had a positive family history and 1486 (44.6%) did not. The polarity of the first episode was available for 3601 (89.2%) of the 4037 patients. Of the 3601 patients, the first episode was depressed in 1748 (48.5%) and manic in 1853 (51.5%).

2.3. Onset location and country median age

This international database has several unique features. Although the data were collected in 36 collection sites in 23 countries, there were 318 unique onset locations (city and country) in 43 countries. Each onset location includes all reported locations within a 1 \times 1 degree grid of latitude and longitude. The number of onset locations from each collection site reflects differences in country size, culture and migration patterns. The number of patients within each onset location varies, and the data within each onset location are correlated [4,5].

There is a large difference in the median age of the population among the countries, varying over 20 years between the oldest (Japan, 45.8 years) and the youngest (South Africa, 25.5 years) [48]. For a disease with a variable age of onset that spans several decades like bipolar disorder, an older age of onset would be expected in a country with an older population [13,27]. Additionally, the country median age, which summarizes the age structure, provides information about the socioeconomic characteristics of a country [48].

2.4. Clustering approach

The clustering analysis was performed in two steps. First, generalized estimating equations (GEE) were used to estimate the effect of the country median age and, in some models the birth cohort, on the age of onset. Second, the residuals from the estimated GEE models, which contain information that was not explained by the GEE model variables, were used for the cluster analysis.

2.5. GEE

All GEE models have the age of onset as the dependent variable. A GEE model was used to accommodate both the correlated data and unbalanced number of patients within the onset locations. All estimates adjust for the correlated onset locations using

clusters, and the country median age as an independent variable. A GEE uses a population averaged or marginal approach, estimating the effect across the entire population rather than within the correlated onset locations [49]. A significance level of 0.01 was used to evaluate estimated coefficients. GEE analyses were performed using geepack 1.1-6 for R.

2.6. Mixture analysis

Mixture analysis was performed using model-based clustering with MCLUST 4.2 for R software [19], as in prior research [24]. Model-based clustering assumes the sample is a mixture of one or more normal distributions, uses a statistical probability model to determine both the number and composition of the clusters, and does not specify in advance the number, shape, volume or orientation of the distributions [17–19]. The best fitting model and number of clusters are selected using the Bayesian Information Criteria (BIC), with the smallest BIC being optimal.

Since the results of this study are population-based, a comparison cannot be directly made with the results for an individual country. However, to confirm the methodology using residuals, a comparison was made using data from just one country. Cluster analysis of age of onset was performed without any adjustments, as in prior studies in a single country [9]. The results were compared to cluster analysis using the age of onset residuals from the GEE model adjusted only for the correlated onset locations within the country. The mean predicted age of onset was added to the cluster midpoint for comparison. As shown in Table 1, there was no difference in the results. Also, the values were similar to prior findings [2,9].

2.7. Impact of birth cohort

A large percentage of 4037 people in this database were born before 1960 (36.8%). As in prior research [14], three birth cohort groups were created: born before 1940, born between 1940 and 1959, and born after 1959. The impact of the birth cohort effect on the clustering was analyzed in three ways. First, using the entire sample, a GEE model was estimated without considering the birth cohort, and cluster analysis was then performed on the residuals. Second, using the entire sample, a GEE model was estimated that also adjusted for the birth cohort, and cluster analysis was then performed on the residuals. Third, a GEE model without the birth cohort adjustment was estimated for only the youngest cohort born after 1959, and cluster analysis was performed on the residuals.

2.8. Clinical variables

The clinical variables of the patients in the subgroups detected by cluster analysis were compared. Clinical variables in this database were family history, gender and polarity of the first episode. The hypomanic and manic data were combined for analysis of polarity. Variables in the subgroups were compared using a Chi2 test. For variables with a significant difference and more than two subgroups, logistic regression models were used for pairwise comparison.

3. Results

Of the 4037 patients, 220 (5.4%) were born before 1940 and had a mean age of onset of 38.4 years, 1267 (31.4%) were born between 1940 and 1959 and had a mean age of onset of 29.5 years, and 2550 (63.2%) were born after 1959 and had a mean age of onset of 22.2 years. The 16.2 years difference between the mean age of onset in the oldest and youngest birth cohort groups influenced the results of the clustering analysis, as shown in Tables 2A–2C. Without considering the birth cohort, the best fitting model

for the entire sample ($n = 4037$) consisted of three normal distributions. The mean age of the three subgroups were 17.24 ± 3.20 , 23.93 ± 5.12 , and 32.20 ± 11.96 years, representing 41.7%, 24.7%, and 33.6% of the sample (Table 2A). With the birth cohort, the best fitting model for the entire sample ($n = 4037$) consisted of two normal distributions. The mean age of two subgroups were 20.7 ± 5.84 and 30.1 ± 10.40 years, representing 62.1% and 37.9% of the sample (Table 2B). Considering only those born after 1959 ($n = 2550$), the best fitting model also consisted of two normal distributions. The mean age of two subgroups were 18.11 ± 3.70 and 25.79 ± 8.41 years, representing 56.9% and 43.1% of the sample (Table 2C).

In all cluster results, more patients in the youngest subgroup had a family history of mood disorders, and a first episode with a depressed polarity (Tables 3A–3C). However, pairwise comparisons of the three subgroups detected without considering the birth cohort, could not distinguish between the middle and oldest subgroups for family history or polarity of first episode. A significant difference in family history and polarity of the first episode was only found when comparing the youngest and middle subgroups, and the youngest and oldest subgroups (Table 4).

4. Discussion

Data in this international study were combined from multiple dissimilar countries, and adjusted for large differences in the country median age. Even with these adjustments, cluster analysis identified three subgroups for the age of onset of bipolar I disorder when the birth cohort is not considered, similar to results from individual countries as summarized by Hamshere et al. [24]. This similarity validates the technique used in this analysis, and, in turn, the cluster analysis on the residuals confirms the presence of subgroups. When adjusting for the birth cohort, or considering only those born after 1959, only two subgroups were found. As in prior studies in which data were unadjusted for the birth cohort, the youngest subgroup was more likely to have a family history of mood disorders [3,22,24], and to have a first episode with a polarity of depression [39,40] when compared to either the middle or older subgroup. However, there was no significant difference between the middle and older subgroups for either family history or polarity of first episode suggesting that the two older subgroups may not be clinically distinct. Since the birth cohort adjustment alters the number of subgroups, the usefulness of this confounder should be investigated in future studies.

The birth cohort effect is a proxy for the cultural environment experienced by different generations of patients and their physicians [43,45,47]. In addition to bipolar disorder, a strong birth cohort effect for age of onset was reported for other psychiatric disorders including depression [12,29,32], schizophrenia [16], substance abuse [28], phobias [36], and symptoms of anxiety [43]. Diverse cultural influences may contribute to the birth cohort effect including the immediate and long-term consequences of World War II [11,31,45,47], stress under totalitarian regimes [6,7], introduction and expansion of psychopharmacology [20,42], evolving diagnostic practices [1,10], changes in societal attitudes to mental illness [21,26,41], changes to family structure and the role of women [32,43], greater exposure to drugs of abuse [12,15,28], and the rise of the information age and social media.

There are several limitations to this study. The data collection process was not standardized across all sites, although diagnosis was based on DSM-IV criteria. Patient reported age of onset is subject to recall or memory bias especially among the elderly [38,46]. The family history data were not validated. Family history data is often inaccurate [25], and may be influenced by cultural attitudes towards mental illness [30]. A genetic anticipation effect may be contributing in part to the birth cohort effect [44]. Ascertainment bias may be present, since patients with bipolar disorder may recognize

symptoms in offspring, resulting in earlier diagnosis. There could also be a selection bias in the age of onset for those born before 1959, since a younger age of onset is associated with a more severe disease course including suicide [40]. This analysis cannot address the importance of the birth cohort effect in any one country. Only three variables were available in this database to evaluate the clinical usefulness of the clustering results. This analysis used the MCLUST mixture algorithm, and clusters determined by other clustering techniques, or by cut-offs based on clinical observation, were not evaluated.

Researchers using mixture analysis have previously noted that a birth cohort effect may influence the composition of the subgroups, or the distribution of some clinical variables within the subgroups [2,9]. Regardless of the cause of the birth cohort effect, ignoring the cohort effect in a statistical analysis of age of onset may produce misleading results.

5. Conclusion

In conclusion, the results of this international study are consistent with prior findings that there are subgroups in the onset of bipolar I disorder [8,9], and that there is a birth cohort effect [14,21]. The birth cohort effect influenced the number and characteristics of the subgroups determined by clustering methodology. Further investigation is needed to determine if including the birth cohort in cluster analysis based on age of onset will identify subgroups that are more useful for clinical research.

Disclosure of interest

The authors declare that they have no conflicts of interest concerning this article.

Acknowledgements

We thank the European College of Neuropsychopharmacology (ECNP) Network Initiative for supporting the European Network of Bipolar Research Expert Centres (ENBREC). This work was also funded in part by the following: Canadian Institutes of Health Research (MA, Grant number 64410); the Research Council of Norway (OAA Grant numbers 213837; 223273; 217776); South-East Norway Health Authority (OAA, Grant number 2013-123); a NHMRC Senior Principal Research Fellowship (M Berk, Grant number 1059660; INSERM (BE, Grant number C0829) and AHPH (BE, Grant number AOR11096); the Spanish Government (AGP, Grant numbers PS09/02002 CIBER Network; EC10-333, PI10/01430, PI10/01746, PI11/01977, PI11/02708, 2011/1064, 11-BI-01, 1677-DJ-030, EC10- 220); European Regional Development Funds (Grant numbers UE/ 2012/FI-STAR, UE/2013/TENDERMH, UE/2013/MASTERMIND), grants from Spanish Government (Grant numbers PI10/01430, PI10/01746, EC10-220, EC10-333, PI11/01977, 20111064, PI11/ 02708, PI12/02077, PI13/02252, PI13/00451), local grants from the Basque Government (Grant numbers 200911147, 2010111170, 2010112009, 2011111110, 2011111113); the Basque Foundation for Health Innovation and Research (Grant number BIO12/AL/002); the Spanish Clinical Research Network (Grant numbers CAI- BER;1392-D-079) and the University of the Basque Country (Grant number IT679-13); Stanley Research Foundation (Grant number 03-RC-003); the Research Council of Norway (IM, Grant numbers ES488722, ES421716); the Regional Health Authority of South Eastern Norway (IM, Grants number 2011085, 2013088); DFG (AR, Grant numbers SFB TRR 58, B06, Z02); the DFG and L nder funds (AR, Grant number RTG1252/2); Medical Research Council of South Africa (DJS); Spanish Ministry of Economy and Competitiveness (EV, Grants number PI12/00912, PN 2008-2011); the Instituto de Salud Carlos III- Subdirecci n General de Evaluaci n y Fomento de la Investigaci n (EV); Fondo Europeo de Desarrollo Regional Uni n Europea. Una manera de hacer Europa

(EV); CIBERSAM (EV); the Comissionat per a Universitats i Recerca del DIUE de la Generalitat de Catalunya to the Bipolar Disorders Group (EV, Grant number 2009 SGR 1022), and the Department of Science and Technology INSPIRE scheme, Government of India (BV).

MB, EA, RA, CB, FB, RB, RHB, TDB, LB, YB, EYWC, MDZ, SD, AF, MAF, KNF, JGF, TG, HH, SH, CH, AI, ETI, FK, SK, BK, RK, MK, BL, RL, CLJ, UL, GM, MM, WM, SM, RM, FGN, CO, YO, AP, DQ, RR, NR, PR, JKR, KS, AMS, ES, CS, SS, AHS, KS, HT, YT, CT, MJW, MZ and PCW have no specific funding to acknowledge.

References

- [1] Akiskal HS. Validating 'hard' and 'soft' phenotypes within the bipolar spectrum: continuity or discontinuity? *J Affect Disord* 2003;73:1–5.
- [2] Azorin JM, Bellivier F, Kaladjan A, Adida M, Belzeaux R, Fakra E, et al. Characteristics and profiles of bipolar I patients according to age at onset: findings from an admixture analysis. *J Affect Disord* 2013;150:993–1000.
- [3] Baldessarini RJ, Tondo L, Vazquez GH, Undurraga J, Bolzani L, Yildiz A, et al. Age at onset versus family history and clinical outcomes in 1665 international bipolar I disorder patients. *World Psychiatry* 2012;11:40–6.
- [4] Bauer M, Glenn T, Alda M, Andreassen OA, Angelopoulos E, Ardaur R, et al. Relationship between sunlight and the age of onset of bipolar disorder: an international multisite study. *J Affect Disord* 2014;167:104–11.
- [5] Bauer M, Glenn T, Alda M, Andreassen OA, Ardaur R, Bellivier F, et al. Impact of sunlight on the age of onset of bipolar disorder. *Bipolar Disord* 2012;14:654–63.
- [6] Bauer M, Priebe S, Kürtén I, Graef KJ, Baumgartner A. Psychological and endocrine abnormalities in refugees from East Germany: part I. Prolonged stress, psychopathology, and hypothalamic-pituitary-thyroid axis activity. *Psychiatry Res* 1994;51:61–73.
- [7] Bauer M, Priebe S, Graef KJ, Kürtén I, Baumgartner A. Psychological and endocrine abnormalities in refugees from East Germany: part II. Serum levels of cortisol, prolactin, luteinizing hormone, follicle stimulating hormone, and testosterone. *Psychiatry Res* 1994;51:75–85.
- [8] Bellivier F, Golmard JL, Henry C, Leboyer M, Schürhoff F. Admixture analysis of age at onset in bipolar I affective disorder. *Arch Gen Psychiatry* 2001;58:510–2.
- [9] Bellivier F, Golmard JL, Rietschel M, Schulze TG, Malafosse A, Preisig M, et al. Age at onset in bipolar I affective disorder: further evidence for three subgroups. *Am J Psychiatry* 2003;160:999–1001.
- [10] Blader JC, Carlson GA. Increased rates of bipolar disorder diagnoses among US child, adolescent, and adult inpatients, 1996–2004. *Biol Psychiatry* 2007;62:107–14.
- [11] Brown AS, van Os J, Driessens C, Hoek HW, Susser ES. Further evidence of relation between prenatal famine and major affective disorder. *Am J Psychiatry* 2000;157:190–5.
- [12] Burke KC, Burke Jr JD, Rae DS, Regier DA. Comparing age at onset of major depression and other psychiatric disorders by birth cohorts in five US community populations. *Arch Gen Psychiatry* 1991;48:789–95.
- [13] Chen WJ, Faraone SV, Orav EJ, Tsuang MT. Estimating age at onset distributions: the bias from prevalent cases and its impact on risk estimation. *Genet Epidemiol* 1993;10:43–59.
- [14] Chengappa KN, Kupfer DJ, Frank E, Houck PR, Grochocinski VJ, Cluss PA, et al. Relationship of birth cohort and early age at onset of illness in a bipolar disorder case registry. *Am J Psychiatry* 2003;160:1636–42.

- [15] da Silva Magalhães PV, Gomes FA, Kunz M, Kapczinski F. Birth cohort and dual diagnosis effects on age at onset in Brazilian patients with bipolar I disorder. *Acta Psychiatr Scand* 2009;120:492–5.
- [16] Di Maggio C, Martinez M, Me´nard JF, Petit M, Thibaut F. Evidence of a cohort effect for age at onset of schizophrenia. *Am J Psychiatry* 2001;158:489–92.
- [17] Fraley C, Raftery AE. How many clusters? Which clustering methods? Answers via model-based cluster analysis. *Comput J* 1998;41:578–88.
- [18] Fraley C, Raftery AE. Model-based clustering, discriminant analysis, and density estimation. *J Am Stat Assoc* 2002;97:611–31.
- [18] Fraley C, Raftery AE. Model-based clustering, discriminant analysis, and density estimation. *J Am Stat Assoc* 2002;97:611–31.
- [19] Fraley C, Raftery AE, Scrucca L. MCLUST: normal mixture modeling for model-based clustering, classification, and density estimation. 2013. R package version 4.2; 2013, <http://www.stat.washington.edu/mclust/>.
- [20] Geddes JR, Miklowitz DJ. Treatment of bipolar disorder. *Lancet* 2013;381:1672–82.
- [21] Gershon ES, Hamovit JH, Guroff JJ, Nurnberger JI. Birth cohort changes in manic and depressive disorders in relatives of bipolar and schizoaffective patients. *Arch Gen Psychiatry* 1987;44:314–9.
- [22] Gonza´lezPintoA, BarbeitoS, Jose´Dí´azF, VegaP, MosqueraF, Lo´pezP, et al. Age at onset in bipolar I disorder: two may be better than three subgroups. *Rev Psiquiatr Salud Ment* 2009;2:29–34.
- [23] Grigoriu-Serbanescu M, Nothen M, Propping P, Poustka F, Magureanu S, Vasilescu R, et al. Clinical evidence for genomic imprinting in bipolar I disorder. *Acta Psychiatr Scand* 1995;92:365–70.
- [24] Hamshere ML, Gordon-Smith K, Forty L, Jones L, Caesar S, Fraser C, et al. Age at onset in bipolar I disorder: mixture analysis of 1369 cases identifies three distinct clinical subgroups. *J Affect Disord* 2009;116:23–9.
- [25] Hardt J, Franke P. Validity, reliability and objectivity of the family history method in psychiatry: a meta-analysis. *Eur Psychiatry* 2007;22:49–58.
- [26] Healy D. The latest mania: selling bipolar disorder. *PLoS Med* 2006;3:e185.
- [27] Heimbuch RC, Matthysse S, Kidd KK. Estimating age of onset distributions for disorders with variable onset. *Am J Hum Genet* 1980;32:564–74.
- [28] Johnson RA, Gerstein DR. Initiation of use of alcohol, cigarettes, marijuana, cocaine, and other substances in US birth cohorts since 1919. *Am J Public Health* 1998;88:27–33.
- [29] JoycePR, Oakley-BrowneMA, WellsJE, BushnellJA, HornblowAR. Birth cohort trends in major depression: increasing rates and earlier onset in New Zealand. *J Affect Disord* 1990;18:83–9.
- [30] KaraszA. Cultural differences in conceptual models of depression. *Soc Sci Med* 2005;60:1625–35.
- [31] Kesternich I, Siflinger B, Smith JP, Winter JK. The effects of world war II on economic and health outcomes across Europe. *Rev Econ Stat* 2014;96:103–18.
- [32] Klerman GL, Weissman MM. Increasing rates of depression. *JAMA* 1989;261:2229–35.
- [33] Kupfer DJ, Frank E, Grochocinski VJ, Cluss PA, Houck PR, Stapf DA. Demographic and clinical characteristics of individuals in a bipolar disorder case registry. *J Clin Psychiatry* 2002;63:120–5.
- [34] Lasch K, Weissman M, Wickramaratne P, Livingston Bruce M. Birth cohort changes in the rates of mania. *Psychiatry Res* 1990;33:31–7.

- [35] Lin PI, McInnis MG, Potash JB, Willour V, MacKinnon DF, DePaulo JR, et al. Clinical correlates and familial aggregation of age at onset in bipolar disorder. *Am J Psychiatry* 2006;163:240–6.
- [36] Magee WJ, Eaton WW, Wittchen HU, McGonagle KA, Kessler RC. Agoraphobia, simple phobia, and social phobia in the National Comorbidity Survey. *Arch Gen Psychiatry* 1996;53:159–68.
- [37] Manchia M, Lampus S, Chillotti C, Sardu C, Ardu R, Severino G, et al. Age at onset in Sardinian bipolar I patients: evidence for three subgroups. *Bipolar Disord* 2008;10:443–6.
- [38] Nivoli AM, Murru A, Pacchiarotti I, Valenti M, Rosa AR, Hidalgo D, et al. Bipolar disorder in the elderly: a cohort study comparing older and younger patients. *Acta Psychiatr Scand* 2014;130:364–73.
- [39] Ortiz A, Bradler K, Slaney C, Garnham J, Ruzickova M, O'Donovan C, et al. An admixture analysis of the age at index episodes in bipolar disorder. *Psychiatry Res* 2011;188:34–9.
- [40] Perlis RH, Miyahara S, Marangell LB, Wisniewski SR, Ostacher M, DelBello MP, et al. Long-term implications of early onset in bipolar disorder: data from the first 1000 participants in the systematic treatment enhancement program for bipolar disorder (STEP-BD). *Biol Psychiatry* 2004;55:875–81.
- [41] Phelan JC, Link BG, Stueve A, Pescosolido BA. Public conceptions of mental illness in 1950 and 1996: what is mental illness and is it to be feared? *J Health Soc Behav* 2000;188–207.
- [42] Schou M. Lithium treatment at 52. *J Affect Disord* 2001;67(1–3):21–32.
- [43] Twenge JM. The age of anxiety? Birth cohort change in anxiety and neuroticism, 1952–1993. *J Pers Soc Psychol* 2000;79:1007–21.
- [44] Visscher PM, Yazdi MH, Jackson AD, Schalling M, Lindblad K, Yuan QP, et al. Genetic survival analysis of age at onset of bipolar disorder: evidence for anticipation or cohort effect in families. *Psychiatr Genet* 2001;11:129–37.
- [45] Wadsworth ME, Kuh DJ. Childhood influences on adult health: a review of recent work from the British 1946 national birth cohort study, the MRC National Survey of Health and Development. *Paediatr Perinat Epidemiol* 1997;11:2–20.
- [46] Warshaw MG, Klerman GL, Lavori PW. Are secular trends in major depression an artifact of recall? *J Psychiatr Res* 1991;25:141–51.
- [47] Willets RC. The cohort effect: insights and explanations. *Br Actuarial J* 2004;10:833–77.
- [48] Central Intelligence Agency. World Factbook 2013–2014.[Accessed 8/30/14] Washington, DC: Central Intelligence Agency; 2013 , <http://www.cia.gov/library/publications/the-world-factbook/index.html>.
- [49] Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics* 1986;42:121–30.

Table 1Comparison of results of cluster analysis of age of onset data for France ($n=371$) using actual data versus residuals.

	Number of subgroups	Youngest subgroup		Middle subgroup		Oldest subgroup	
		n (%)	Mean (SD)	n (%)	Mean (SD)	n (%)	Mean (SD)
Actual data	3	156 (42.0)	17.03 (2.45)	145 (39.1)	24.71 (5.02)	70 (18.9)	36.52 (11.26)
Residuals ^a	3	156 (42.0)	-7.94 (2.45)	145 (39.1)	-0.26 (5.02)	70 (18.9)	11.55 (11.26)
Residuals mean + overall mean age of onset ^b	3	156 (42.0)	17.03 (2.45)	145 (39.1)	24.71 (5.02)	70 (18.9)	36.52 (11.26)

The entries in bold demonstrate that using the actual data, and the residuals mean + overall mean age of onset, produced the same result.

^a Residuals calculated using generalized estimating equation (GEE) estimate of age of onset as a function of a constant with 28 onset locations within France.^b The overall mean of the estimated GEE age of onset is 24.97 years.**Table 2A**Results of cluster analysis of age of onset data for all patients without birth cohort ($n=4037$)^a.

Number of subgroups	Youngest subgroup		Middle subgroup		Oldest subgroup	
	n (%)	Mean (SD)	n (%)	Mean (SD)	n (%)	Mean (SD)
3	1685 (41.7)	17.24 (3.20)	995 (24.7)	23.93 (5.12)	1357 (33.6)	32.20 (11.96)

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant and the country median age with 318 onset locations. The overall mean of the estimated GEE age of onset is 25.38 years.**Table 2B**Results of cluster analysis of age of onset data for all patients with birth cohort ($n=4037$)^a.

Number of subgroups	Youngest subgroup		Oldest subgroup	
	n (%)	Mean (SD)	n (%)	Mean (SD)
2	2506 (62.1)	20.7 (5.84)	1531 (37.9)	30.1 (10.40)

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant, the country median age and birth cohort group with 318 onset locations. The overall mean of the estimated GEE age of onset is 25.40 years.**Table 2C**Results of cluster analysis of age of onset data for patients born after 1959 ($n=2550$)^a.

Number of subgroups	Youngest subgroup		Oldest subgroup	
	n (%)	Mean (SD)	n (%)	Mean (SD)
2	1452 (56.9)	18.11 (3.70)	1098 (43.1)	25.79 (8.41)

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant and the country median age with 263 onset locations. The overall mean of the estimated GEE age of onset is 22.22 years.

Table 3A

Patient characteristics in subgroups from cluster analysis of age of onset data without birth cohort^a.

	Youngest subgroup		Middle subgroup		Oldest subgroup		Chi ²
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Gender (<i>n</i> =4037)							<i>P</i> =0.935
Female	989	58.7	590	59.3	795	58.6	
Male	696	41.3	405	40.7	562	41.4	
Family history (<i>n</i> =3334)							<i>P</i> <0.001
Yes	819	61.9	446	52.5	583	50.2	
No	504	38.1	404	47.5	578	49.8	
Polarity of first episode (<i>n</i> =3601)							<i>P</i> <0.001
Depressed	790	54.1	388	42.6	570	46.4	
Manic	671	45.9	523	57.4	659	53.6	

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant and the country median age with 318 onset locations.

Table 3B

Patient characteristics in subgroups from cluster analysis of age of onset data with birth cohort^a.

	Youngest subgroup		Oldest subgroup		Chi ²
	<i>n</i>	%	<i>n</i>	%	
Gender (<i>n</i> =4037)					<i>P</i> =0.853
Female	1477	58.9	879	58.6	
Male	1029	41.1	634	41.4	
Family history (<i>n</i> =3334)					<i>P</i> <0.001
Yes	1200	59.7	648	49.0	
No	811	40.3	675	51.0	
Polarity of first episode (<i>n</i> =3601)					<i>P</i> <0.001
Depressed	1135	51.4	613	44.0	
Manic	1072	48.6	781	56.0	

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant, the country median age and birth cohort group with 318 onset locations

Table 3C
 Patient characteristics in subgroups from cluster analysis of age of onset data for patients born after 1959^a.

	Youngest sub-group		Oldest subgroup		Chi ²
	<i>n</i>	%	<i>n</i>	%	
Gender (<i>n</i> = 2550)					<i>P</i> = 0.641
Female	845	58.2	628	57.2	
Male	607	41.8	470	42.8	
Family history (<i>n</i> = 2091)					<i>P</i> < 0.001
Yes	686	59.3	457	48.9	
No	471	40.7	477	51.1	
Polarity of first episode (<i>n</i> = 2272)					<i>P</i> < 0.001
Depressed	665	52.2	400	40.0	
Manic	608	47.8	599	60.0	

^a Modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant and the country median age with 263 onset locations.

Table 4
 Pairwise comparison of patient characteristics within subgroups from cluster analysis without birth cohort^a.

	Youngest vs. middle				Youngest vs. oldest				Middle vs. oldest			
	OR ^c	<i>P</i>	95% CI ^b		OR ^c	<i>P</i>	95% CI ^b		OR ^c	<i>P</i>	95% CI ^b	
			2.5%	97.5%			2.5%	97.5%			2.5%	97.5%
Polarity of first episode (<i>n</i> = 3601)												
Manic (<i>n</i> = 1853)	1.587	< 0.001	1.344	1.876	1.361	< 0.001	1.169	1.585	0.858	0.081	0.722	1.019
Depressed (<i>n</i> = 1748) ^d												
Family history (<i>n</i> = 3334)												
Yes (<i>n</i> = 1848)	0.679	< 0.001	0.571	0.809	0.621	< 0.001	0.529	0.728	0.914	0.320	0.765	1.091
No (1486) ^d												

^a Subgroups modeled using residuals from generalized estimating equation (GEE) estimate of age of onset as a function of a constant and the country median age with 318 onset locations. Pairwise comparison using logistic regression.

^b Confidence interval.

^c Odds ratio.

^d Reference category.